



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07D 501/22		A1	(11) International Publication Number: WO 98/45299 (43) International Publication Date: 15 October 1998 (15.10.98)
<p>(21) International Application Number: PCT/EP98/01953</p> <p>(22) International Filing Date: 2 April 1998 (02.04.98)</p> <p>(30) Priority Data: A 570/97 4 April 1997 (04.04.97) AT </p> <p>(71) Applicant (<i>for all designated States except US</i>): BIOCHEMIE GESELLSCHAFT MBH [AT/AT]; A-6250 Kundl (AT).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): STURM, Hubert [AT/AT]; Leopoldstrasse 40, A-6020 Innsbruck (AT). WOLF, Siegfried [AT/AT]; Bruggerstrasse 4, A-6230 Brixlegg (AT). LUDESCHER, Johannes [AT/AT]; Kleinsoell 101, A-6252 Breitenbach (AT).</p> <p>(74) Agent: BECKER, Konrad; Novartis AG, Patent- und Markenabteilung, Lichtstrasse 35, CH-4002 Basel (CH).</p>			<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: CRYSTALLINE AMINE SALT OF CEF DINIR</p> <p>(57) Abstract</p> <p>Cefdinir in the form of a salt with dicyclohexylamine, a process for its production and its use in the purification of impure cefdinir.</p>			


FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		

CRYSTALLINE AMINE SALT OF CEFIDINIR

The present invention relates to intermediates in the purification of cefdinir, i.e. 7-(Z)-[2-(2-aminothiazol-4-yl)-2-hydroxyiminoacetamido]-3-vinyl-3-cephem-4-carboxylic acid of formula



Cefdinir may be used, e.g. in form of a monohydrate, as a pharmaceutical, e.g. antibiotic; see e.g. Y.Inamoto, Toshiyuki Chiba, Toshiaki Kiamimura und Takao Takaya, J.Antibiotics Vol XLI, No 6, 829, (1988).

Cefdinir may be obtained in impure form according to known production processes. It was now surprisingly found that impure cefdinir may be purified via the formation of a salt, e.g. in crystalline form, thereof.

In one aspect the present invention provides a compound of formula I in the form of a salt, e.g. crystalline, with dicyclohexylamine.

A compound of formula I in the form of a salt with dicyclohexylamine may be produced as follows:

Cefdinir, e.g. in the form of a solvate, such as a hydrate, e.g. in impure form, e.g. as obtained in a production process of cefdinir, such as a mixture of cefdinir and impurities, e.g. such as a mixture of by-products originating from the production process of cefdinir and cefdinir; may be treated in the presence of a solvent, e.g. in dissolved or suspended form, with dicyclohexylamine. A solvent includes any solvent which is inert towards cefdinir or towards cefdinir in the form of a salt with dicyclohexylamine, e.g. a polar organic solvent, such as amides, e.g. dimethylformamide; alcohols, e.g. methanol oder ethanol; ketones, e.g acetone; e.g. in combination with water and water. A solvent system, e.g. mixtures of individual solvents, e.g. as described above may be used. A preferred solvent system may be e.g. acetone/water, including e.g. a ratio of about 100:1 such as 50:1, e.g. 20:1 to 1:5; such as 10:1 to 1:3, e.g. 5:1 to 2:1, e.g. about 1:1. Per equivalent of cefdinir about one equivalent or more, such as 5; e.g. 3, such as 2 equivalents of dicyclohexylamine may be used, e.g. combined

with the mixture of impure cefdinir in a solvent. A compound of formula I in the form of a salt with dicyclohexylamine may crystallize e.g. from a reaction solution, or, e.g. a suspension of a compound of formula I in a solvent may be converted into a crystal suspension of a compound of formula I in the form of a salt with dicyclohexylamine. An anti-solvent may be added to the reaction mixture, e.g. in order to complete crystallization. An anti-solvent includes solvents wherein a compound of formula I in the form of a salt with dicyclohexylamine is insoluble or soluble only to a small extent if added to the solution or suspension of a compound of formula I in the form of a dicyclohexylamine, e.g. apolar solvents; e.g. ethers, such as diethylether, tetrahydrofuran; or a ketone, e.g. acetone. A compound of formula I in the form of a salt with dicyclohexylamine may be isolated from the reaction mixture, e.g. as conventional, e.g. by filtration, centrifugation.

A compound of formula I in the form of a salt with dicyclohexylamine may be obtained in pure form, e.g. in 98% purity and more, such as 99% to 100% purity; e.g. the amount of impurities present in cefdinir in impure form used for salt formation may be decreased; e.g. impurities of 10% and more in cefdinir in impure form may be decreased to impurities of 1% and less, e.g. 0 to 1% in cefdinir in the form of a salt with dicyclohexylamine.

A compound of formula I in the form of a salt with dicyclohexylamine may be further purified by re-suspension or re-dissolution as described above, e.g. in an (anti) solvent (system), e.g. as described above.

In another aspect the present invention provides a process for the production of a compound of formula I in the form of a salt with dicyclohexylamine, e.g. in crystalline form, comprising treating a compound of formula I, e.g. in form of a solvate, such as a hydrate, in a solvent with dicyclohexylamine and isolating a compound of formula I in the form of a salt with dicyclohexylamine, e.g. in crystalline form.

Cefdinir in free form, e.g. in the form of a solvate, such as a hydrate, e.g. monohydrate and in purified form, e.g. in respect with impure cefdinir used for the formation of a salt of a compound of formula I with dicyclohexylamine, may be obtained from a compound of formula I in the form of a salt with dicyclohexylamine, e.g. as conventional for setting free a compound which is in the form of a salt, e.g. in the form of an amine salt; e.g. by adjusting an appropriate pH, e.g. 1.5 to 4, such as 2 to 3; of a mixture, e.g. a solution, of cefdinir in the form of a salt with dicyclohexylamine with a solvent, e.g. in the presence of water, preferably in water, e.g. by addition of an acidic agent, such as an organic or inorganic acid, preferably

an inorganic acid, e.g. sulphuric acid. Cefdinir, e.g. in the form of hydrate, e.g. monohydrate may crystallize and may be isolated, e.g. as conventional, e.g. by filtration, centrifugation. A compound of formula I may be obtained according to the process of the present invention as such or in the form of a solvate, e.g. a hydrate, e.g. a monohydrate. A compound of formula I obtained according to the process of the present invention as such may be converted into a compound of formula I in the form of a solvate, e.g. a hydrate, such as a monohydrate and vice versa.

In another aspect the present invention provides a process for the production of a compound of formula I, e.g. in form of a solvate, such as a hydrate, e.g. monohydrate comprising converting a compound of formula I in the form of a salt with dicyclohexylamine, e.g. in crystalline form, into a compound of formula I, e.g. in the form of a solvate, and isolating a compound of formula I, e.g. in the form of a solvate.

In another aspect the present invention provides a process for the purification of cefdinir in a mixture of a compound of formula I with impurities, comprising forming a salt of a compound of formula I with dicyclohexylamine; and converting a compound of formula I in the form of a salt with dicyclohexylamine, e.g. in crystalline form, into a compound of formula I, e.g. in the form of a solvate, and isolating a compound of formula I, e.g. in the form of a solvate.

A compound of formula I in the form of a salt with dicyclohexylamine is useful in the purification of cefdinir in impure form.


In another aspect the present invention provides the use of a compound of formula I in the form of a salt with dicyclohexylamine, e.g. in crystalline form in the purification of a mixture of a compound of formula I with impurities.

The present invention has several surprising advantages:

A compound of formula I in the form of a salt with dicyclohexylamine may be in acrystalline form; cefdinir in the form of a salt may be obtained in surprising high purity, e.g. 98% purity and more, e.g. 98% to 100%;- production of the salt is simple; cefdinir obtained from the salt may be surprisingly pure, e.g. 98% and more, e.g. 99% to 100%.

It is surprising that cefdinir under the basic conditions of the salt formation according to the present invention is stable, because from e.g. Yoshihiko Okamoto et al., J. of Pharmaceutical Sciences, Vol 85, No 9, 976, (1996) it is known that cefdinir may be instable in a basic environment; it was e.g. found that cefdinir in the presence of other amines, e.g. tert.-octylamine under similar conditions may be, e.g. heavily, degraded.

In another aspect the present invention provides crystalline 7-(Z)-[2-(2-aminothiazol-4-yl)-2-hydroxyiminoacetamido]-3-vinyl-3-cephem-4-carboxylic acid.dicyclohexylammonium salt of formula



In the following examples, which illustrate the invention more fully but do in no way limit its scope, all temperatures are given in degrees Celsius. Purity of a compound obtained is determined by HPLC.

Example 1

7-(Z)-[2-(2-Aminothiazol-4-yl)-2-hydroxyiminoacetamido]-3-vinyl-3-cephem-4-carboxylic acid in form of a salt with dicyclohexylamine

10 g of crude cefdinir, e.g. as obtained in a cefdinir production process, in 50 ml of water and 50 ml of acetone are treated under stirring with 5 ml of dicyclohexylamine. A solution is obtained and 7-(Z)-[2-(2-Aminothiazol-4-yl)-2-hydroxyiminoacetamido]-3-vinyl-3-cephem-4-carboxylic acid in form of a salt with dicyclohexylamine crystallizes. 250 ml of acetone are added to the crystall suspension which is stirred for ca. 30 minutes at room temperature. The crystals are filtrated off, washed with acetone and dried. Crystalline 7-(Z)-[2-(2-Aminothiazol-4-yl)-2-hydroxyiminoacetamido]-3-vinyl-3-cephem-4-carboxylic acid in form of a salt with dicyclohexylamine in a purity of 98.6 % is obtained. Mp: 175° (decomposition).

¹H-NMR (DMSO-d₆): 9.41 (d, 1H, J = 8.1 Hz, NH); 7.12 (s, 2H, NH₂); 6.99 (dd, 1H, J = 11.4 and 17.7 Hz, CH = CH₂); 6.64 (s, 1H, thiazol); 5.60 (dd, 1H, J = 4.8 and 8.1 Hz, H₇); 5.15 (d, 1H, J = 17.7 Hz, CH = CH₂); 5.04 (d, 1H, J = 4.8 Hz, H₆); 4.94 (d, 1H, J = 11.4 Hz, CH = CH₂); 3.52, 3.39 (AB d, 1H, J = 17 Hz, H₂); 3.21 (m, 2H); 2.05 (m, 4H); 1.8 (m, 4H); 1.6 (m, 2H); 1.2 - 1.4 (m, 10H).

Example 2

7-(Z)-[2-(2-Aminothiazol-4-yl)-2-hydroxyiminoacetamido]-3-vinyl-3-cephem-4-carboxylic acid in the form of a monohydrate

10 g of 7-(Z)-[2-(2-Aminothiazol-4-yl)-2-hydroxyiminoacetamido]-3-vinyl-3-cephem-4-carboxylic acid in form of a salt with dicyclohexylamine, obtained according to Example 1 are dissolved in 175 ml of water at a temperature of ca. 35-40° and treated with active charcoal. Active charcoal is filtrated off and the pH of the solution obtained is adjusted to pH 2.5 by addition of 5ml of sulphuric acid at ca. 35°. 7-(Z)-[2-(2-Aminothiazol-4-yl)-2-hydroxyiminoacetamido]-3-vinyl-3-cephem-4-carboxylic acid in the form of a monohydrate precipitates, is filtrated off, washed with water and dried. 7-(Z)-[2-(2-Aminothiazol-4-yl)-2-hydroxyiminoacetamido]-3-vinyl-3-cephem-4-carboxylic acid in the form of a monohydrate in a purity of 99% is obtained.


Example 3**Production of crude cefdinir**

40 g of 7-amino-3-vinyl-3-cephem-4-carboxylic acid in 400 ml of dichloromethane are treated with 55.7 ml of N,O-bistrimethyl-silylacetamid. The mixture obtained is stirred for ca. 2 hours at room temperature, cooled to 0° and treated with 52.2 g of 2-(Z)-(2-aminothiazol-4-

yl)-2-acetoxyiminoacetic acid chloride in the form of a hydrochloride in small portions. The mixture obtained is stirred for ca. 90 minutes at 0° and added under stirring to a mixture of 44.55 g of NaHCO₃, 600 ml of water und 100 ml of dichloromethane of a temperature of 5°. The pH of the mixture obtained is adjusted to a pH of ca. 7.2 - 7.3 with a saturated aqueous solution of NaHCO₃. The phases formed are separated. To the aqueous phase 300 ml of water are added and 28.7 g of NH₄Cl. The pH of the mixture obtained is adjusted to pH 8 by addition of an aqueous 10% K₂CO₃ solution and the mixture is stirred for ca. 80 minutes. A solution is obtained. The pH of the solution obtained is adjusted to pH 3 by addition of 5M sulphuric acid. 7-(Z)-[2-(2-amino-thiazol-4-yl)-2-hydroxyiminoacetamido]-3-vinyl-3-cephem-4-carboxylic acid precipitates, is filtrated off, washed with water and dried. 7-(Z)-[2-(2-amino-thiazol-4-yl)-2-hydroxyiminoacetamido]-3-vinyl-3-cephem-4-carboxylic acid in a purity of 94.3 % is obtained.

Patent claims


1. A compound of formula



in the form of a salt with dicyclohexylamine.

2. A process for the production of a compound of formula I as defined in claim 1 in the form of a salt with dicyclohexylamine comprising treating a compound of formula I in a solvent with dicyclohexylamine and isolating a compound of formula I in the form of a salt with dicyclohexylamine.
3. A process for the production of a compound of formula I as defined in claim 1 comprising converting a compound of formula I in the form of a salt with dicyclohexylamine into a compound of formula I and isolating a compound of formula I.
4. A process for the purification of a compound of formula I as defined in claims 1 in a mixture of a compound of formula I with impurities, comprising forming a salt of a compound of formula I with dicyclohexylamine; and converting a compound of formula I in the form of a salt with dicyclohexylamine into a compound of formula I and isolating a compound of formula I.
5. A process according to any one of claims 3 to 4 comprising converting a compound of formula I in the form of a salt with dicyclohexylamine into a compound of formula I in the form of a solvate.
6. A process according to any one of claims 3 to 5 comprising isolating a compound of formula I in the form of a solvate.
7. Use of a compound of claim 1 in the purification of a mixture of a compound of formula I with impurities.

8. A compound of formula I as defined in claim I in the form of a salt with dicyclohexylamine according to any one of claims 1 to 7, which is in crystalline form.
9. Crystalline 7-(Z)-[2-(2-aminothiazol-4-yl)-2-hydroxyiminoacetamido]-3-vinyl-3-cephem-4-carboxylic acid.dicyclohexylammonium salt of formula



INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/01953

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D501/22

According to International Patent Classification(IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 559 334 A (T. TAKAYA ET AL.) 17 December 1985 see column 2, line 5; claims ---	1,8,9
Y	WO 97 07121 A (BIOCHEMIE GMBH) 27 February 1997 see page 11 - page 12; claims ---	1-9
Y	GB 1 038 529 A (FUJISAWA PHARMACEUTICAL CO., LTD.) 10 August 1966 see page 10, example 32; claim 33 ---	1-9
A	EP 0 304 019 A (FUJISAWA PHARMACEUTICAL CO., LTD.) 22 February 1989 see claims ---	1-9
		-/-

Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

° Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

10 August 1998

Date of mailing of the international search report

18/08/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Chouly, J

INTERNATIONAL SEARCH REPORT

atational Application No

PCT/EP 98/01953

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P , A	WO 97 24358 A (HANMI PHARMACEUTICAL CO., LTD.) 10 July 1997 see claims -----	1-9

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 98/01953

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
US 4559334	A 17-12-1985	AT 385994	B	10-06-1988
		AU 576735	B	08-09-1988
		AU 1927783	A	05-04-1984
		CA 1206956	A	01-07-1986
		CH 657857	A	30-09-1986
		DK 427083	A, B,	31-03-1984
		EP 0105459	A	18-04-1984
		FI 833370	A, B,	31-03-1984
		FR 2533926	A	06-04-1984
		GB 2127812	A, B	18-04-1984
		JP 1926846	C	25-04-1995
		JP 6057713	B	03-08-1994
		JP 62294687	A	22-12-1987
		PT 77426	B	27-02-1986
		SU 1309911	A	07-05-1987
WO 9707121	A 27-02-1997	AT 403049	B	27-10-1997
		AT 136995	A	15-03-1997
		AU 6821396	A	12-03-1997
		EP 0844999	A	03-06-1998
GB 1038529	A	DE 1545794	A	11-12-1969
		FR 5564	M	02-01-1968
		SE 329165	B	05-10-1970
EP 304019	A 22-02-1989	AT 123221	T	15-06-1995
		AU 617347	B	28-11-1991
		CA 1297096	A	10-03-1992
		DE 3853901	D	06-07-1995
		DE 3853901	T	12-10-1995
		ES 2072856	T	01-08-1995
		HK 18496	A	09-02-1996
		IE 67348	B	20-03-1996
		JP 1250384	A	05-10-1989
		JP 1943842	C	23-06-1995
		JP 6074276	B	21-09-1994
		KR 9708126	B	21-05-1997
		MX 9203468	A	01-09-1992
		US 4935507	A	19-06-1990

INTERNATIONAL SEARCH REPORT

Information on patent family members

ational Application No

PCT/EP 98/01953

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9724358	A 10-07-1997	NONE	